arXiv:cond-mat/0110396v1 19 Oct 2001

Comment on "Peculiar Scaling of Self-Avoiding Walk Contacts"

Baiesi, Orlandini and Stella have recently shown [1] that the set of all non-bonded nearest neighbour (NN) contacts between the two halves of a self-avoiding walk is a multifractal characterized by a universal exponent, τ. In the presence of NN interactions, and in two dimensions, they have shown that $\tau \sim 1.93$ above the θ temperature, T_θ, wheras $\tau = 1$ at and below T_θ. They have shown further that the exponent τ could be superuniversal with a value, $\tau = 2$, for contacts between a self-avoiding walk and a confining wall. The purpose of this comment is to demonstrate that such contact-scaling is broken for the recently proposed Interacting Growth Walk (IGW) [2].

IGW starts from an arbitrarily chosen site, $\mathbf{r}_0$, of a regular d-dimensional lattice of coordination number z, whose sites are initially 'unoccupied' (by monomers). The first step is made in one of the z available directions, by choosing an 'unoccupied' NN of $\mathbf{r}_0$, say $\mathbf{r}_1$, at random with equal probability. Let $\{\mathbf{r}_j^m \mid m = 1, 2, ..., z_j\}$ be the 'unoccupied' NNs available for the j^{th} step of the walk. If $z_j = 0$, the walk can not grow further because it is geometrically 'trapped'. It is, therefore, discarded and a fresh walk is started from $\mathbf{r}_0$. If $z_j \neq 0$, the walk proceeds by choosing one of the available sites, say $\mathbf{r}_j^m$, with a probability proportional to $exp[-\beta n_{NN}^m(j)\epsilon_0]$, where $n_{NN}^m(j)$ is the number of non-bonded NN sites of $\mathbf{r}_j^m$ and $\beta = 1/k_BT$, k_B is the Boltzmann constant. At 'infinite' temperature ($\beta = 0$), the walk generated will be the same as the Kinetic Growth Walk (KGW), reference given in [2]. However, at finite temperatures, the walk will prefer to step into a site with more non-bonded NN contacts when the energy per contact, ϵ_0, is negative. We may set $\epsilon_0 = -1$ without loss of generality.

At any given temperature, we make a large number of attempts [3] to generate N-step IGWs, identify the non-bonded NN contacts between the two halves of the walks, locate their centres of mass and compute the ensemble average of the q-th power of the average radius of gyration, $< R_c^q(N) > \sim N^{\sigma_q}$, ($q > 0$). From the observed power-law dependence of the moments, we obtained σ_q as a function of q for IGWs on a square lattice, and have plotted them in the upper inset of Fig.1 for $\beta = 0, 1/2, 1, 2$ and 300. The fact that σ_q/q depends on q for all these temperatures is related to the specific nature of the corresponding distributions, $P_\beta(R_c, N)$, which we have shown in the lower inset of Fig.1 for $N = 300$. It is clear that the distribution has a power-law form, $P_{\beta=0}(R_c, N = 300) \sim R_c^{-\tau}$, only at infinite temperature. By demanding that the q-th moment, obtained from the scaling form, $P_{\beta=0}(R_c, N) \sim R_c^{-\tau} f(R_c/N^\nu)$ where the end-to-end distance exponent $\nu = 3/4$ for the KGW, be consistent with the monte carlo data in Fig.1, we get the contact-exponent, $\tau \sim 1.90 \pm 0.02$. That this value of τ is consistent with the assumed scaling form is further confirmed in Fig.1, where we have shown that the distributions corresponding to $N = 200$ and $N = 300$ could be collapsed reasonably well. Thus, the IGW at infinite temperature (or equivalently, the KGW) has the same contact-scaling behaviour as a self-avoiding walk However, at finite temperatures, it has none.



FIG. 1. Data collapse for $N = 200$ (filled circle) and $N = 300$ (star) at infinite temperature ($\beta = 0$) with the contact-exponent $\tau = 1.9$. Upper Inset: σ_q as a function of q for $\beta = 0, 0.5, 1, 2$ and 300. Lower Inset: The distributions, $P_\beta(R_c, N = 300)$, for $\beta = 0, 0.5, 1, 2$ and 300 from left to right.

S.L. Narasimhan[1*], P.S.R. Krishna[1*], K.P.N.Murthy[2] and M. Ramanadham[1]
[1]Solid State Physics Division, Bhabha Atomic Research Centre, Mumbai - 400 085, India.
[2]Materials Science Division, Indira Gandhi Centre for Atomic Research, Kalpakkam - 603 102, India.
[*]glass@apsara.barc.ernet.in

---

[1] M. Baiesi, E. Orlandini and A. L. Stella, Phys. Rev. Lett. **87**,070602 (2001).

[2] S. L. Narasimhan, P. S. R. Krishna, K. P. N. Murthy and M. Ramanadham, preprint *cond-mat*/0108097, and references therein; submitted to Phys. Rev. E.

[3] Typically ∼ 200 million attempts are made to generate walks upto a maximum of $N = 500$ on a square lattice; sample attrition is the most severe problem for $\beta = 0$ and it becomes less and less severe as the value of β increases.